

Chemical Compositions and Antioxidant Activities of The Essential Oils of Some Medicinal and Aromatic Plants

Sibel Soycan Öneç^{1*}, Zümrüt Açıkgoz², Figen Kırkpınar², Tuncay Küme³,
Çiğdem Şeremet Tuğalay², Özer Hakan Bayraktar²

¹Namık Kemal Üniversitesi Ziraat Fakültesi Zootečni Bölümü, ²Ege Üniversitesi Ziraat Fakültesi Zootečni Bölümü,

³Dokuz Eylül Üniversitesi Tıp Fakültesi Biyokimya ve Klinik Biyokimya Bölümü, İzmir

*İletişim (correspondence): e-posta: ssonenc@nku.edu.tr; Tel:+90 (282) 250 2186; Fax: +90 (282) 250 250 9929

Gönderim tarihi (Received): 08 Aralık 2015; Kabul tarihi (Accepted): 05 Şubat 2016

Abstract

The present study was conducted to determine the chemical compositions and antioxidant activities of some essential oils of some medicinal and aromatic plants widely used in feed or food industry. The major compounds of the essential oils of cinnamon, cumin, laurel, mint, oregano, rosemary and sage are cinnamaldehyde propylene glycol acetal (41.50%), cuminaldehyde (44.01%), 1,8 cineole (39.55), (+) pullegon (67.80%), carvacrol (59.03%), 1,8 cineole (30.12%) and (+) camphor (17.15%), respectively. There were significant differences in the antioxidant activities of these essential oils ($P<0.01$). In terms of the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay, laurel essential oil (79.00%) demonstrated the highest antioxidant activity, followed by that from cumin (75.98%), oregano (75.81%), mint (69.49%), sage (69.01%), cinnamon (68.83%) and rosemary (63.88%). To conclude, the DPPH free radical scavenging activities of all essential oils from some medicinal and aromatic plant species are significantly greater than those of vitamin E and Trolox ($P<0.01$).

Keywords: Essential oil, chemical composition, antioxidant activity, aromatic plants, spices.

Bazı Tıbbi ve Aromatik Bitkilerden Elde Edilen Uçucu Yağların Kimyasal Kompozisyonları ve Antioksidan Aktiviteleri

Öz

Bu çalışma yem ve gıda endüstrisinde yaygın olarak kullanılan bazı uçucu yağların kimyasal kompozisyonlarının ve antioksidan aktivitelerinin belirlenmesi amacıyla yürütülmüştür. Tarçın, kimyon, defne, nane, kekik, biberiye ve adaçayı uçucu yağlarının başlıca bileşenleri sırasıyla cinnamaldehyde propilene glycol acetat (%41.50), cuminaldehyde (%44.01), 1,8 cineole (%39.55), (+) pullegon (%67.80), carvacrol (%59.03), 1,8 cineole (%30.12) ve (+) camphor (%17.15)'dur. Bu uçucu yağların antioksidan aktivitelerinde önemli düzeyde farklılıklar bulunmuştur ($P<0.01$). DPPH yöntemine göre en yüksek antioksidan aktiviteyi defne uçucu yağı (%79.00) göstermiştir. Bunu, kimyon (%75.98), kekik (%75.81), nane (%69.49), ada çayı (%69.01), tarçın (%68.83) ve biberiye (%63.88) uçucu yağları izlemiştir. Sonuç olarak, bazı tıbbi ve aromatik bitkilerden elde edilen uçucu yağların DPPH serbest radikal yakalama aktivitelerinin Vitamin E ve Trolox'dan önemli düzeyde yüksek olduğu belirlenmiştir ($P<0.01$).

Anahtar kelimeler: Uçucu yağ, kimyasal kompozisyon, antioksidan aktivite, DPPH

Introduction

In feed or food industry, synthetic antioxidants such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), *tert*-butylhydroquinone (TBHQ) and propyl gallate (PG) have been widely used to prevent or delay chemical deterioration caused by lipid oxidation. These additives are preferred due to their low cost and high stability over many years. However, consumers are now concerned about the health risks attributed to some of these synthetic antioxidants (namely, BHT and BHA) (Prasad et al., 2009). Thus, natural antioxidants are in high demand because of their

health-enhancing and disease-risk-preventing properties (Su et al., 2007).

Due to their antiviral, antibacterial, antifungal and antioxidant properties, plant essential oils are multifunctional (García et al., 2003; Matan et al., 2006; Sukutta et al., 2008; Reichling et al., 2009). Globally, essential oils are increasingly used as natural antioxidants that limit oxidative degradation of lipids. The antioxidant activity of plant essential oils might be related to the presence of hydroxyl groups on their phenolic compounds (Shahidi and Nacz, 2004). The effectiveness of phenolic compounds in retarding lipid oxidation is mainly due to their free radical scavenging

activity, transition metal-chelating potential and/or singlet oxygen-quenching capacity (Shahidi and Naczki, 2004; Carmona-Jimenez et al., 2014).

In recent years, several *in vitro* and *in vivo* studies have been conducted to determine the antioxidant effects of aromatic plants and spices. Extensive *in vitro* studies on different rosemary, oregano, sage and cumin have been performed (Stefanovits-Bányai et al., 2003; Embuscado 2015). Due to the use of plant extracts in most of these studies, information about the antioxidant activities of plant essential oils is limited.

Various *in vitro* methods have been used to measure and compare the antioxidant activities of aromatic plants and spices. 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical scavenging activity is a rapid, simple and inexpensive method and can be applicable to both solid and liquid samples. In addition, this method is not specific to any particular antioxidant component and measures the overall antioxidant capacity of the samples (Carmona-Jimenez et al., 2014).

The present study was conducted to determine the chemical composition and antioxidant activities of cinnamon, cumin, laurel, mint, oregano, rosemary and sage essential oils. The DPPH free radical scavenging activities of these oils were also compared to those of vitamin E and trolox, which were used as reference antioxidants.

Material and Method

Material

In this study, essential oils obtained from cinnamon (*Cinnamomum zeylanicum* L.), cumin (*Cuminum cyminum* L.), laurel (*Laurus nobilis* L.), mint (*Mentha pulegium* L.), oregano (*Origanum onites* L.), rosemary (*Rosmarinus officinalis* L.) and sage (*Salvia triloba* L.) were used. These essential oils were obtained from Ege Lokman Botanical Plant Industry Trade Ltd. Company (Kırkağaç, Manisa, Turkey). The herbs used for steam-distilled essential oils were detailed in Table 1.

Method

Tris hydroxymethyl-aminomethane was purchased from Riedel-de Haën (Steinheim, Germany). Hydrochloric acid (HCl), absolute ethanol, Tween-20 and DPPH (2,2-diphenyl-1-picrylhydrazyl) were obtained from Sigma-Aldrich (Steinheim, Germany). Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) was obtained from Fluka Chemie AG (Steinheim, Germany). Water was purified using the Easy Pure RF UV Small Batch Water System (Barnstead, USA).

Components of the essential oil were separated on an HP Agilent 6890 gas chromatograph coupled to an Agilent 5973 mass spectrometer. An HP-Agilent (Nr:19091N-116 Innowax) capillary column (60 m x 0.32 mm, 0.25 µm film thickness) was used directly into the ion source of the MSD. The carrier gas was helium at a constant pressure of 9.05 psi. The oven programming was conducted with an initial temperature of 70°C, a rate of 7°C/minute, a final temperature of 210°C and an injection volume of 1 µL.

In the present study, the chemical compositions of the essential oils of cinnamon, cumin, laurel, mint, oregano, rosemary and sage were identified using gas chromatography–mass spectrometry (GC/MS, HP 6890 GC/5973 MSD) at the Ege University Centre R & D and Pharmacokinetic Applications–Environmental & Food Analysis Laboratories–Food Control Laboratory (Bornova, İzmir, Turkey) according to the United States Pharmacopeia and the National Formulary (USP, 1995).

The free radical scavenging activities of seven essential oils, vitamin E and trolox were determined in accordance with the procedures recommended by Sacchetti et al. (2005). This method involves measuring the scavenging ability of free radicals. This spectrophotometric assay uses the stable radical DPPH as a reagent. DPPH was added into the tubes for analysis as follows: 900 µL of buffer (100 mM Tris-HCl, pH 7.4), 40 µL of solvent (ethanol), 50 µL of emulsifying agent (Tween-20, 0.5%) and 1000 µL of

Table 1. Botanical characteristics of some medicinal and aromatic plants used in the study.

| Herbs | Family | Botanical name | Plant part | Collection site |
|----------|-----------|-------------------------------|----------------------------|-----------------|
| Cinnamon | Lauraceae | <i>Cinnamomum zeylanicum</i> | Cortex cinnamomi zeylanici | Seylan |
| Cumin | Apiaceae | <i>Cuminum cyminum</i> | Fructus cumini cymini | Aksehir |
| Laurel | Lauraceae | <i>Laurus nobilis</i> | Folium lauri nobili | Hatay |
| Mint | Lamiaceae | <i>Mentha pulegium</i> | Folium pulegi | Mugla |
| Oregano | Lamiaceae | <i>Origanum onites</i> | Folium origani | Izmir |
| Rosemary | Lamiaceae | <i>Rosmarinus officinalis</i> | Folium rosmarini | Mersin |
| Sage | Lamiaceae | <i>Salvia triloba</i> | Folium salviae trilobae | Izmir |

colour compound (0.5 mM DPPH). Next, 10 µL of type 1 pure water (resistivity > 0.1 megohm and conductivity <1 µS) was added into the blank tube, 50 µL of trolox was added into the trolox tube, and 10 µL of each essential oil was added into the test tubes. After 70 minutes of incubation in the dark at room temperature, the absorbance at 517 nm was measured against the blank with a visible spectrophotometer. Vitamin E and trolox were used as positive controls. The values are presented as the mean of five analyses. The radical-scavenging activities of the samples, expressed as per cent inhibition of DPPH ($I\%$) were calculated from the absorbance of the blank (A_B) and of the sample (A_A) by following the equation from Carmona-Jimenez (2014).

$$I\% = [(A_B - A_A) / A_B] \times 100$$

where A_B and A_A are the absorbance values of the blank and essential oil solutions checked after 70 minutes, respectively.

The results are expressed as mean \pm standard error (SE) (n=5). Free radical scavenging activities of different essential oils were analysed by one-way ANOVA using the General Linear Model Procedure of SAS (1989). Means were compared using and Dunnett's means test at the 5% significance level.

Results and Discussion

Chemical composition of essential oils

The chemical compositions of the essential oils of cinnamon, cumin, laurel, mint, oregano, rosemary and sage are presented in Table 2.

In our study, it was shown that the major compounds in the essential oil of cinnamon are cinnamaldehyde propylene glycol, cinnamaldehyde and propylene, which amounted to 41.50%, 35.28% and 2.76% of the contents, respectively. Tomaino et al. (2005) determined that the essential oil of cinnamon contains cinnamic aldehyde (67.9%), eugenol (6.72%) and β -caryophyllene (5.03%) at room temperature. In another study, Ünlü et al. (2010) found out that the essential oil from *C. zeylanicum* bark contains (E)-cinnamaldehyde (68.95%), benzaldehyde (9.94%), (E)-cinnamyl acetate (7.44%), limonene (4.42%) and eugenol (2.77%). The level of cinnamaldehyde reported in the present study is lower than that found by Ünlü et al. (2010).

The major compounds in the essential oil of cumin were 44.01% cuminaldehyde, 23.85% *p*-cymene, 8.25% safranal and 8.11% β -pinene. Gachkar et al. (2007) identified the major compounds of the essential oil of *C.*

cyminum as α -pinene, limonene, 1,8-cineole and linalool (29.1%, 21.5%, 17.9% and 10.4 %, respectively). The cuminaldehyde content of the essential oil of cumin defined in the present study is lower than reported by Soyca-Onenç and Akkan (2009).

In Table 2, it was shown that the essential oil of laurel is especially rich in 1,8-cineole (39.55%), α -terpinyl acetate (21.72%) and terpinen-4-ol (4.38%). Sangun et al. (2007) determined that the essential oil of *L. nobilis* leaves contains 1,8-cineole (46.61-59.94%), α -terpinyl acetate (11.94-25.70%) and terpinen-4-ol (1.82-2.20%). The current level of 1,8-cineole in laurel was lower than that of Sangun et al. (2007) while α -terpinyl acetate content was in line what Sangun et al. (2007) determined. However, the present terpinen-4-ol content was higher than that of Sangun et al. (2007).

In our study, it was found that the essential oil of mint contains (+)-pulegone (67.80%), isomenthone (15.77%), piperitenone (3.91%), piperitone oxide (1.44%) and menthol (1.43%) as the main compounds. The pulegone content of *M. Pulegium* was reported by Stoyanova et al. (2005) as 42.9-45.4% and by El-Ghorab (2006) as 43.5% and piperitone content of the latter study 12.2% as the major compounds. The pulegone content in mint essential oil determined in the present study is higher than that found by Stoyanova et al. (2005) and El-Ghorab (2006). The level of piperitone was lower than that reported by El-Ghorab (2006).

Tomaino et al. (2005) reported that the major components in oregano oil (*Origanum floribundum* Munby) kept at room temperature are carvacrol (48.9%), thymol (5.03%) and *p*-cymene (11.77%). The major components of the essential oil of oregano in this study were carvacrol, thymol, and *p*-cymene (59.03%, 12.04% and 6.37%) respectively. The carvacrol content reported in this study is higher than that reported by Tomaino et al. (2005).

The major components of the essential oil of rosemary were characterised by Gachkar et al. (2007) as piperitone, α -pinene, linalool, 1,8-cineole, camphor, borneol, camphene and bornyl acetate; and by Verma et al. (2010) as camphor, α -pinene, 1,8-cineole, borneol, verbenone, linalool, limonene, *exo*-bornyl acetate, terpinen-4-ol and camphene. Soyca-Onenç and Akkan (2009) reported that the major components in rosemary essential oil are 1,8-cineole (15-30 and 14.34%), camphor (5-10 and 23.54%) and borneol (10-20 and 26.16%). In our study, the essential oil of rosemary

Table 2. Chemical composition of the essential oils obtained from some medicinal and aromatic plants.

| Cinnamon | | Cumin | | Laurel | | Mint | |
|-------------------------|-------|-----------------------|-------|-----------------------------|--------|-----------------------|-------|
| Compounds | % | Compounds | % | Compounds | % | Compounds | % |
| Cinnamaldehyde | | Cuminaldehyde | 44.01 | 1,8 Cineole | 39.55 | (+) Pulegon | 67.80 |
| propylene glycol acetal | 41.50 | | | | | | |
| Cinnamaldehyde | 35.28 | p- Cymene | 23.85 | α - Terpinyl acetate | 21.72 | Isomenthon | 15.77 |
| Propylene glycol | 2.76 | Safranal | 8.25 | Terpinen-4-ol | 4.38 | Piperitenone | 3.91 |
| Unidentified | 19.66 | β - Pinene | 8.11 | β -Phellandrene | 3.70 | Piperitone Oxide | 1.44 |
| | | γ -Terpinene | 2.11 | p-Cymene | 3.43 | Menthol | 1.43 |
| | | Carotol | 2.00 | α - Pinene | 3.16 | α - Pinene | 1.05 |
| | | (-) α - Cedren | 1.93 | Eugenol metyl ether | 3.01 | β - Pinene | 1.00 |
| | | Carvacrol | 1.86 | β - Pinene | 2.80 | Piperitone | 0.77 |
| | | Anethol | 1.65 | α -Terpineole | 2.05 | Neomenthol | 0.74 |
| | | Cuminic Alcohol | 1.52 | Linalool | 1.99 | Limonen | 0.73 |
| | | α - Thujen | 0.75 | Euganol | 1.91 | Amyl Vinyl Carbinol | 0.55 |
| | | Limonen | 0.58 | Limonen | 1.63 | Spathulenol | 0.49 |
| | | α - Terpineole | 0.57 | Gamma Terpinen | 1.62 | α -Terpineole | 0.34 |
| | | p- Cymene-8- ol | 0.41 | Cis α -Bisabolene | 1.09 | β -Phellandrene | 0.33 |
| | | Others | 1.24 | Others | 5.80 | Others | 0.73 |
| | | Unidentified | 1.15 | Unidentified | 2.16 | Unidentified | 2.91 |
| Total | 100 | | 100 | | 100 | | 100 |
| Oregano | | Rosemary | | Sage | | | |
| Compounds | % | Compounds | % | Compounds | % | | |
| Carvacrol | 59.03 | 1,8 Cineole | 30.12 | (+) Camphor | 17.15 | | |
| Thymol | 12.04 | α - Pinene | 12.80 | 1,8 Cineole | 14.84 | | |
| p-Cymene | 6.37 | (+)Campher | 12.75 | β -Thujon | 8.01 | | |
| Γ - Terpinen | 3.86 | (+) Borneol | 6.38 | Caryophyllene | 6.53 | | |
| Linalool | 2.73 | Caryophyllene | 5.48 | α - Humulene | 5.98 | | |
| (+) Borneol | 2.09 | Camphene | 4.06 | (+) Borneol | 5.34 | | |
| Terpinen-4-ol | 1.83 | Terpineole | 3.99 | α - Pinene | 5.01 | | |
| B-Caryophyllene | 1.67 | p-Cymene | 2.81 | Carvacrol | 3.71 | | |
| B- Bisabolene | 1.58 | (+)-Limonen | 2.33 | Camphene | 3.60 | | |
| A-Terpinen | 1.24 | L-Bornylester | 1.79 | Viridiflorol | 3.37 | | |
| Sabinen | 1.03 | Carvacrol | 1.61 | L-Bornylester | 3.26 | | |
| α - Humulene | 0.83 | Linalool | 1.49 | α - Thujon | 3.10 | | |
| α - Thujon | 0.74 | β - Pinene | 1.16 | α - Pinene | 2.20 | | |
| α - Terpineol | 0.65 | Myrcen | 1.04 | p-Cymene | 2.12 | | |
| Others | 4.47 | Others | 8.42 | Others | 10.957 | | |
| Unidentified | 0.32 | Unidentified | 3.43 | Unidentified | 4.77 | | |
| Total | 100 | | 100 | | 100 | | |

mainly comprised 1,8-cineole (30.12%), α -pinene (12.80%), (+) camphor (12.75%), (+) borneol (6.38%) and caryophyllene (5.48%), respectively. Similar to our findings, Fu et al. (2007) reported that the major components of the essential oil of this plant are 1,8-cineole (48.5 and 27.23%), α -pinene (15.4 and 19.43%) and camphor (10 and 14.26%). As shown in Table 2, (+)

camphor, 1,8-cineole, β -thujone, caryophyllene, α -humulene, (+) borneol and α -pinene are the major compounds in the essential oil of sage, at 17.15, 14.84, 8.01, 6.53, 5.98, 5.34 and 5.01%, respectively. Soycan-Onenç and Akkan (2009) reported that the major components of the essential oil of *S. triloba* are 1,8-cineole (34.40%), camphor (18.40%), camphene

(8.80%), α -pinene (7.90%), borneol, α -terpinyl acetate (4.80%) and β -pinene (4.40%).

Antioxidant activity of essential oils and vitamin E

The DPPH free radical scavenging activities in the essential oils of cinnamon, cumin, laurel, mint, oregano, rosemary and sage are shown in Table 3.

Table 3. DPPH-free radical scavenging activities of the essential oils and vitamin E ($X \pm SE$).

| The source of essential oils | Inhibition, % |
|--------------------------------------|-------------------------------|
| Medicinal and aromatic plants | |
| Sage | 69.01 \pm 0.30 ^c |
| Rosemary | 63.88 \pm 1.29 ^d |
| Laurel | 79.00 \pm 0.54 ^a |
| Oregano | 75.81 \pm 1.65 ^b |
| Cumin | 75.98 \pm 1.16 ^b |
| Mint | 69.49 \pm 1.47 ^c |
| Cinnamon | 68.83 \pm 0.88 ^c |
| Control substances | |
| Vitami E | 22.08 \pm 0.85 ^f |
| Trolox | 28.79 \pm 0.57 ^e |
| Probablity (<i>P value</i>) | 0.0001 |

^{a-f}: Means within a column with different superscripts differ significantly ($P < 0.05$).

There were significant differences in the free radical scavenging activities between essential oils investigated in this study ($P < 0.01$). Among the investigated essential oils, laurel had the highest free radical scavenging activity at 79 \pm 0.54%, while rosemary essential oil showed the lowest free radical scavenging activity at 63.88 \pm 1.29%. The difference in inhibition between the essential oil of cumin (75.98 \pm 1.16%) and that of oregano (75.81 \pm 1.65%) was not statistically significant. A similar situation was found for the essential oils of mint, sage and cinnamon; inhibition levels of these essential oils were 69.49 \pm 1.47%, 69.01 \pm 0.30% and 68.83 \pm 0.88%, respectively. The free radical scavenging activities of trolox and vitamin E were determined to be 28.79 \pm 0.57% and 22.08 \pm 0.85%.

In this study, the highest DPPH free scavenging activity was found for the essential oil of laurel, which contains approximately 40% 1,8-cineole. In accord with our findings, Politeo et al. (2006) also found that the essential oil of *L. nobilis* (with a major component of 1,8-cineole) showed strong antioxidant activity.

The essential oil of cumin (75.98 \pm 1.16%) demonstrated the next highest antioxidant activity (Table 3). Lu et al. (2011) found that *C. cyminum* extract has low antioxidant capacity (18.12%). This result was probably

due to the low total phenolic content of the cumin extract used (9.00 \pm 0.15 mg GAE/g DW) in their study.

While the essential oil of oregano had similar DPPH inhibition levels to the essential oil of cumin, showing significantly higher antioxidant activity than those of mint, sage, cinnamon and rosemary.

DPPH inhibition of the essential oil of mint was significantly higher than that of the essential oil of rosemary only (Table 3). In a previous study, Mata et al. (2007) determined that water extracts of *M. spicata* and *M. pulegium* (IC₅₀=5.7 and 8.9 μ g/ml, respectively) have higher DPPH radical scavenging activities than the ethanol extracts of these species (IC₅₀=65.2 and 24.9 μ g/ml, respectively), the water and ethanol extracts of *R. officinalis* (IC₅₀=37.3 and 36.0 μ g/ml respectively) and BHT (IC₅₀=15.7 μ g/ml). Kamkar et al. (2010) found that water and methanol extracts from *M. pulegium* have similar free radical scavenging activities to BHT but higher free radical scavenging activities than the essential oil of *M. pulegium*. These researchers noted that the lower antioxidant activity in the essential oil of *M. pulegium* could be due to the lower rate of presence of different antioxidants in the essential oil.

In our study, it was found that the essential oil of sage had a significantly higher DPPH inhibition level than the essential oil of rosemary. Dorman et al. (2003) determined that there was no difference in the DPPH radical scavenging activities between the aqueous extracts of *R. officinalis* and *S. officinalis* (IC₅₀=236.5 μ g/ml and 265.8 μ g/ml, respectively). These plants' extracts showed greater anti-radical effects than that from the aqueous extract of *O. vulgare* (IC₅₀=335 μ g/ml).

This study showed that the highest anti-radical activity was in the essential oil of laurel, which contains approximately 40% 1,8-cineole. Many essential oils isolated from various plant species belonging to different genera contain relatively high amounts of oxygenated monoterpenes such as borneol, borneol acetate, camphor, carvone, 1,8-cineole, linalool, linalool acetate, limonene oxide, menthol, menthone, terpinen-4-ol, α -terpineol (Kotan et al. 2007), which has demonstrated poor ability of inhibiting oxidation (Cherrat et al. 2014). This high antioxidant activity in the essential oil of laurel can be attributed to the presence of monoterpene hydrocarbons, which was the α -terpinyl acetate (21.72%).

Prasad et al. (2009) reported that the DPPH radical scavenging activities of cinnamon leaf extracts from

five species increased with increasing concentration, and the highest DPPH radical scavenging activity was observed in *C. zeylanica* (92.1%) at 100 µg/ml. Moreover, in the studies carried out by Muchuweti et al. (2007), Dudonné et al. (2009) and Lu et al. (2011) *C. zeylanica* extract showed a high antioxidant capacity, with mean values of 84.43, 92 and 87.45%, respectively, for DPPH inhibition. It was hypothesised that the DPPH radical scavenging activities of cinnamon species may result from the hydrogen-donating ability of phenols and flavonoids (Prasad et al., 2009). The DPPH inhibition levels for cinnamon reported by these authors are greater than we determined in our study. The discrepancy in the results may be due to the use of extracts instead of essential oils in these four studies. In addition, the type of extracting solvent might affect the antioxidant activities of spices and herbs, as reported by Su et al. (2007).

Among the examined essential oils, it was determined that the lowest antioxidant capacity was found for the essential oil of rosemary (Table 3). In an earlier study carried out by Sacchetti et al. (2005), it was observed that the DPPH free radical scavenging activity of the essential oil of *R. officinalis* was lower than that of the essential oil of *T. vulgaris* and approximately two times higher than that of trolox. The inhibition values for the essential oil of rosemary essential oil were similar to each other.

The major components of the essential oil of rosemary (1,8-cineole, camphor, camphene, borneol, borneol acetate, linalool, limonene, α -pinene, piperitone, verbenone) were different from those of rosemary extracts (carnosol, carnosic acid, rosmanol, rosmarinic acid, naringin, hispidulin, cirsimaritin, caffeic acid, vanillic acid, apigenin) (Luis and Johnson, 2005). Carnosol, carnosic acid and rosmarinic acid were the most active antioxidant components (Wei and Ho, 2006), while 1,8-cineol, α -pinene, camphor and verbenone are the most active antimicrobial components (Moghtader and Afzali, 2009) in rosemary. In our study, the essential oil of rosemary rich in 1,8-cineole (30.12%) content exhibited the lowest antioxidant activity, which supports the above statements.

Conclusion

In this study, chemical compositions and *in vitro* antioxidant activities of essential oils in some medicinal and aromatic plants widely used in food or feed industry were investigated. The DPPH radical scavenging activities of the essential oils were ranked as followings:

laurel > cumin \approx oregano > mint \approx sage \approx cinnamon > rosemary. It is important that these results could be checked by further *in vivo* studies, especially for the feed industry, in terms of defining the availability of essential oils.

Acknowledgement

This study was funded by the Scientific and Technological Research Council of Turkey (Project number TOVAG-106 O 621).

References

- Carmona-Jimenez Y, Garcia-Moreno MV, Igartuburu JM, Barroso CG. 2014. Simplification of the DPPH assay for estimating the antioxidant activity of wine and by-products. *Food Chemistry* 165:198-204.
- Cherrat L, Espina L, Bakkali M, Pagan R, Laglaoui A. 2014. Chemical composition, antioxidant and antimicrobial properties of *Mentha pulegium*, *lavandula stoechas* and *Satureja calamintha* Scheele essential oils and evaluation of their bactericidal effect in combined processes. *Innovative Food Science and Emerging Technologies* 22:221-229.
- Dorman HJD, Peltoketo A, Hiltunen R, Tikkanen MJ. 2003. Characterisation of the antioxidant properties of de-odourised aqueous extracts from selected Lamiaceae 200 herbs. *Food Chemistry* 83:255-262.
- Dudonné S, Vitrac X, Coutière P, Woillez M, Mérillon J-M. 2009. Comparative study of antioxidant properties and total phenolic content of 30 plant extracts of industrial interest using DPPH, ABTS, FRAP, Sod, and ORAC assays. *Journal of Agricultural and Food Chemistry* 57:1768-1774.
- El-Ghorab AH. 2006. The chemical composition of *Mentha pulegium* L. essential oil from Egypt and its antioxidant activity. *Journal of Essential Oil Bearing Plants* 9:183-195.
- Embuscado ME. 2015. Spices and herbs: Natural sources of antioxidants-a mini review. *Journal of Functional Foods* 18:811-819. doi:10.1016/j.jff.2015.03.005.
- Fu Y, Zu Y, Chen L, Shi X, Wang Z, Sun S, Efferth T. 2007. Antimicrobial activity of clove and rosemary essential oils alone and in combination. *Phytotherapy Research* 21:989-994.
- Gachkar L, Yadegari D, Rezaei MB, Taghizadeh M, Astaneh SA, Rasooli I. 2007. Chemical and biological characteristics of *Cuminum cyminum* and *Rosmarinus officinalis* essential oils. *Food Chemistry* 102:898-904.
- García CC, Talarico L, Almeida N, Colombres S, Duschatzky C, Damonte EB. 2003. Virucidal activity of essential oils from aromatic plants of San Luis, Argentina. *Phytotherapy Research* 17:1073-1075.

- Kamkar A, Javan AJ, Asadi F, Kamalinejad M. 2010. The antioxidative effect of Iranian *Mentha pulegium* extracts and essential oil in sunflower. *Food and Chemical Toxicology* 48:1796-1800.
- Kotan R, Kordali S, Çakır A. 2007. Screening of antibacterial activities of twenty-one oxygenated monoterpenes. *Zeitschrift für Naturforschung C* 62:507-513.
- Lu M, Yuan B, Zeng M, Chen J. 2011. Antioxidant capacity and major phenolic compounds of spices commonly consumed in China. *Food Research International* 44:530-536.
- Luis JC, Johnson CB. 2005. Seasonal variations of rosmarinic and carnosic acids in rosemary extracts. Analysis of their in vitro antiradical activity. *Spanish Journal of Agricultural Research* 3(1):106-112.
- Mata AT, Proença C, Ferreira AR, Serralheiro MLM, Nogueira JMF, Araújo MEM. 2007. Antioxidant and antiacetylcholinesterase activities of five plants used as Portuguese food spices. *Food Chemistry* 103:778-786.
- Matan N, Rimkeeree H, Mawson AJ, Chompreeda P, Haruthaithansan V, Parker M. 2006. Antimicrobial activity of cinnamon and clove oils under modified atmosphere conditions. *International Journal of Food Microbiology* 107:180-185.
- Moghtader M, Afzali D. 2009. Study of the antimicrobial properties of the essential oil of rosemary. *American-Eurasian Journal of Agricultural & Environmental Sciences* 5(3):393-397.
- Muchuweti M, Kativu E, Mupure CH, Chidewe C, Ndhlala AR, Benhura MAN. 2007. Phenolic composition and antioxidant properties of some spices. *American Journal of Food Technology* 2(5):414-420.
- Prasad KN, Yang B, Dong X, Jiang G, Zhang H, Xie H, Jiang Y. 2009. Flavonoid contents and antioxidant activities from *Cinnamomum* species. *Innovative Food Science and Emerging Technologies* 10:627-632.
- Politeo O, Jukić M, Miloš M. 2006. Chemical composition and antioxidant activity of essential oils of twelve spice plants. *Croatica Chemica Acta* 79(4):545-552.
- Reichling J, Schnitzler P, Suschke U, Saller R. 2009. Essential oils of aromatic plants with antibacterial, antifungal, antiviral, and cytotoxic properties-An Overview. *Forschende Komplementärmedizin* 16:79-90.
- SAS, 1989. SAS/STAT User's Guide AS/STAT User's Guide (Release 6.03 Edition) SAS Institute Inc, Cary, NC, USA.
- Sacchetti G, Maietti S, Muzzoli M, Scaglianti M, Manfredini S, Radice M, Bruni R. 2005. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chemistry* 91:621-632.
- Sangun MK, Aydın E, Timur M, Karadeniz H, Çalışkan M, Özkan A. 2007. Comparison of chemical composition of the essential oil of *Laurus nobilis* L. Leaves and fruits from different regions of Hatay, Turkey. *Journal of Environmental Biology* 28(4):731-733.
- Shahidi F, Naczki M. 2004. Phenolics in food and nutraceuticals. CRC Press LLC, 2000 N.W. Corporate Blvd., Boca Raton, Florida 33431. Print ISBN 1-58716-138-9.
- Stefanovits-Bányai É, Tulok MH, Hegedüs A, Renner C, Varga IS. 2003. Antioxidant effect of various rosemary (*Rosmarinus officinalis* L.) clones. *Acta Biologica Szegediensis* 47:11-113.
- Su L, Yin J-J, Charles D, Zhou K, Moore J, Yu L. 2007. Total phenolic contents, chelating capacities, and radical-scavenging properties of black peppercorn, nutmeg, rosehip, cinnamon and oregano leaf. *Food Chemistry* 100: 990-997.
- Sukutta U, Haruthaithansan V, Chantarapanont W, Dilokkunanant U, Suppakul P. 2008. Antifungal activity of clove and cinnamon oil and their synergistic against postharvest decay fungi of grape in vitro. *Kasetsart Journal (Natural Science)* 42:169-174.
- Soycan-Onenç S, Akkan S. 2009. The effects of some aromatic plants on rumen volatile fatty acids. V. National Animal Nutrition Congress 30 September-3 October Çorlu-Tekirdağ, p. 202-207.
- Stoyanova A, Georgie V, Kula J, Majda T. 2005. Chemical composition of the essential oil of *Mentha pulegium* from Bulgaria. *Journal of Essential Oil Research* 17:475-477.
- USP, 1995. The U.S. Pharmacopeia National Formulary. USP 23 NF 18, p.1755.
- Tomaino A, Cimino F, Zimbalatti V, Venuti V, Sulfaro V, De Pasquale A, Saija A. 2005. Influence of heating on antioxidant activity and the chemical composition of some spice essential oils. *Food Chemistry* 89:549-554.
- Ünlü M, Ergene E, Vardar-Ünlü G, Sivas-Zeytinoğlu H, Vural N. 2010. Composition, antimicrobial activity and in vitro cytotoxicity of essential oil from *Cinnamomum zeylanicum* Blume (*Lauraceae*). *Food Chemistry Toxicology* 48:3274-3280.
- Verma RS, Sashidhara KV, Yadav A. 2010. Essential oil composition of 'blue flower rosemary' (*Rosmarinus officinalis* L.) from subtropical India. *Acta Pharmaceutica Scientia* 52:427-430.

Wei G-J, Ho C-T. 2006. A stable quinone identified in the reaction of carnosol, a major antioxidant in rosemary, with 2,2-diphenyl-1-picrylhydrazyl radical. *Food Chemistry* 96:471-476.